	rood Anergy Action Flan	
ALLERGY	TO:	Place Child's
Student's	D.O.B:Teacher:	Picture Here
Asthmatic Y	Yes* No *High risk for severe reaction	
♦ SIGNS	OF AN ALLERGIC REACTION ◆	
Systems:	Symptoms:	
•MOUTH •THROAT* •SKIN •GUT •LUNG* •HEART*	itching & swelling of the lips, tongue, or mouth itching and/or a sense of tightness in the throat, hoarseness, and hacking cough hives, itchy rash, and/or swelling about the face or extremities nausea, abdominal cramps, vomiting, and/or diarrhea shortness of breath, repetitive coughing, and/or wheezing "thready" pulse, "passing-out"	h
The severity threatening	of symptoms can quickly change. *All above symptoms can potentially prosituation.	ogress to a life-
♦ ACTIO	N FOR MINOR REACTION ♦	
1. If only sy	mptom(s) are:	lose/route
Then call:		
2. Mother 3. Dr	, Father, or e	mergency contacts.
If condition d	loes not improve within 10 minutes, follow steps for Major Reaction below.	
♦ ACTIO	N FOR MAJOR REACTION •	
1. If ingestio	on is suspected and/or symptom(s) are:	
give	medication/dose/route	IMMEDIATELY!
Then call:		
3. Mother	uad (ask for advanced life support), Father, or e	emergency contacts.
	DO NOT HESITATE TO CALL RESCUE SQUAD!	
Parent's Signa	atureDate Doctor's Signature	Date

Food Alloway Action Dlan


EMERGENCY CONTACTS	TRAINED STAFF MEMBERS	
1.	1Room _	
Relation:Phone: 2.	2Room _	
Relation:Phone:3.	3Room _	
Relation:Phone:		

EPIPEN® AND EPIPEN® JR. DIRECTIONS

1. Pull off gray activation cap.



2. Hold black tip near outer thigh (always apply to thigh).



3. Swing and jab firmly into outer thigh until Auto-Injector mechanism functions. Hold in place and count to 10. The EpiPen® unit should then be removed and taken with you to the Emergency Room. Massage the injection area for 10 seconds.

For children with multiple food allergies, use one form for each food.

